

A Small Dose of Arsenic
Or
An Introduction to the Health Effects of Arsenic

By Steven G. Gilbert, PhD, DABT



From Wikipedia - Elemental arsenic — mineral specimen
<https://en.wikipedia.org/wiki/Arsenic>

Dossier

Name: Arsenic

Use: wood preservative, pesticides, semiconductor manufacturing

Source: coal combustion, drinking water, environment, medical drug, seafood

Recommended daily intake: none (not essential)

Absorption: inhalation, intestine – inorganic high, organic low, skin

Sensitive individuals: children

Toxicity/symptoms: Peripheral nervous system (tingling in hands in feet), skin cancer (ingestion), lung cancer (inhalation); Hyperpigmentation (keratosis) of palms and soles; vascular complications

Regulatory facts: EPA - Drinking water 10 $\mu\text{g/L}$ (0.01 ppm, 10 ppb)

EPA – RfD – 0.3 $\mu\text{g/kg/day}$

OSHA - Workplace air 10 $\mu\text{g/m}^3$

ATSDR - MRL - 0.3 $\mu\text{g/kg/day}$

General facts: long history of use as medicine and poison

Environmental: global environmental contaminate, bioaccumulates in fish and shellfish (mostly in a form that is not harmful)

Recommendations: avoid, do not use arsenic treated lumber, test drinking water

Case Studies

(Henry Adams)...he found himself invariably taking for granted, as a political instinct, with out waiting further experiment—as he took for granted that arsenic poisoned—the rule that a friend in power is a friend lost.

Henry Adams (1838–1918). *The Education of Henry Adams*. 1918

Arsenic in Drinking Water

Arsenic in drinking water is a worldwide problem affecting the lives of millions of people. High levels of arsenic in local soil or rock contaminate the local water supply. In the United States, the federal government has struggled for many years to establish standards of arsenic in the drinking water. The U.S. Environmental Protection Agency has recently lowered the standard from 50 ppm (50 $\mu\text{g/L}$) to 10 ppm. This standard will require additional treatment of a number of municipal water supplies particularly in the western United States (see map in presentation). The standard is being lowered because chronic exposure to low levels of arsenic can cause skin cancer and others illnesses. Even at the new standard of 10 ppm, there is a risk of cancer.

In other areas of the world, such as Bangladesh, elevated arsenic levels in the drinking water is more acutely life threatening. People were encouraged to establish local wells to reduce exposure to bacteria-contaminated drinking water. It was subsequently discovered that many of these wells have high levels of arsenic in the water. It is estimated that 75 million people of Bangladesh are exposed to arsenic-contaminated water that will result in 200,000 to 270,000 deaths from cancer each year. In addition, people suffer from skin changes on the palms of hands and soles of feet. (for additional information see presentation).

Pressure-treated wood

By far the largest use of arsenic is in treating wood to prevent decay or insect damage. Several compounds are used, but the vast majority of wood is treated with a pesticide called chromated copper arsenate (CCA), first used in the 1940s. CCA is a water-based mixture of inorganic salts of chromium, copper, and arsenic that is forced into the wood under pressure. Wood treated with CCA is still found in decks, playground equipment, outdoor furniture, fences, construction lumber, utility poles, piers, and pilings. The amount of arsenic in treated wood can be quite large. A standard eight-foot length of treated 2" x 4" lumber contains as much as 15 grams of arsenic. To put this in perspective the lethal dose of arsenic in humans is 70 to 200 mg or about 1 mg/kg. Since December

31, 2003, CCA was no longer used in wood for most residential settings, including decks and play sets. There are a number of arsenic-free wood preservatives on the market that are registered use in treated wood for domestic use.

The health risks of exposure to arsenic-treated lumber have been debated for years, although it is well known that inhaling sawdust from construction with treated lumber can be quite dangerous. Ideally the arsenic-based wood preservative is “fixed” to the wood, but research has shown that arsenic leaches from the wood with rainfall and that arsenic can be rubbed off from the surface by hand contact. Arsenic contamination of soil under decks often exceeds hazardous waste cleanup standards. Children who play on decks or other treated surfaces pick up arsenic on their hands and later ingest some of the arsenic when they put their hands in their mouth or pick up food. Health professionals, the wood preserving industry, and public interest groups have hotly debated the hazards of these exposures. In 2002, producers of arsenic-treated wood reached an agreement with EPA to phase out the residential uses of arsenic treated lumber, including decks, play equipment, fences, etc. CCA will still be available for commercial uses such as utility poles. The alternative wood treatment most used to replace CCA is a copper-based preservative called ammoniacal copper quaternary, or ACQ. ACQ has a much lower toxicity to humans than CCA.

Introduction and History

I pray my companion, if he wishes for bread, to ask me for bread, and if he wishes for sassafras or arsenic, to ask me for them, and not to hold out his plate, as if I knew already.

Ralph Waldo Emerson (1803–1882).

People long ago recognized that depending on the dose, arsenic could either treat an illness or be used as a poison to cause death. Its medicinal use to treat syphilis and amebic dysentery ended with the introduction of penicillin and other antibiotics in the twentieth century. Arsenic-based compounds are currently used to treat some forms of cancer. As a poison, arsenic trioxide (As_2O_3) has several desirable qualities: it looks like sugar, it is tasteless, and it only takes about a tenth of gram to kill

As Arsenic

Atomic Number: 33

Atomic Mass 74.92

someone. While its use as a human poison has greatly declined, arsenic is still used as a pesticide particularly in growing cotton, as an herbicide and as a wood preservative.

Arsenic poisoning from well water remains a serious world wide human health concern. Internationally, in West Bengal and Bangladesh more than 75 million people are exposed to arsenic-laden water that threatens their health. People of Argentina, Chile, and Taiwan also have elevated arsenic in their drinking water. In the United States, federal agencies fiercely debate arsenic drinking water standards, which would limit the amount of arsenic in municipal wells. This is particularly relevant to areas of the western United States that have elevated arsenic their drinking water.

Arsenic is a versatile metal, forming various compounds, either inorganic or organic, with a complex chemistry. Inorganic arsenic is widely distributed in nature usually in the trivalent form (As^{3+}) but also as pentavalent arsenic (As^{5+}). The trivalent forms include arsenic trioxide, sodium arsenite, and arsenic trichloride. Organic arsenic, much less toxic than inorganic arsenic, is produced in a biomethylation process by many organisms including humans and shellfish.

Arsenic use and production has declined with recognition of its toxicity and the development of suitable replacements. It is not mined but produced as by product of smelting for copper, lead and zinc. The last U.S. smelter producing arsenic in Tacoma, Washington closed in 1985. Smelters typically released the trivalent arsenic trioxide and lead into the atmosphere which contaminated the local environment leaving an unwelcome legacy for local residents. This use of arsenic is also being phased out, such as its use as a wood preservative in CCA, as reflected by a decline in imported arsenic from 20,000 metric tons in 2002 and 2003 to less than 8,000 metric tons in 2007. Arsenic is used in the manufacture of silicon-based computer chip technology and in glass manufacture to control color. Inorganic arsenic is no longer used as a pesticide in cotton fields and orchards, but some forms of organic arsenic continue to be applied to cotton fields. Inorganic arsenic is also released from coal-fired electric generation facilities, and cigarette smokers inhale some arsenic from tobacco. Organic arsenic compounds are also used as a feed additive to enhance growth of poultry and swine.

We are exposed to constant but low levels of arsenic, unless receiving greater exposure in an occupational setting or from arsenic-contaminated drinking water. Normally, the background air contains less than $0.1 \mu\text{g}/\text{m}^3$ and drinking water less than $5 \mu\text{g}/\text{L}$, but water levels can be significantly higher. Food usually supplies less than $10 \mu\text{g}/\text{day}$ of arsenic but can be higher with the consumption of fish and particularly shellfish, which can have arsenic levels up to $30 \mu\text{g}/\text{g}$. The majority of arsenic in food is organic, a form that is generally less toxic than inorganic arsenic. The total average daily exposure to arsenic is about $20 \mu\text{g}/\text{day}$ from food and water (assuming 2000 mL/day average water consumption at $5 \mu\text{g}/\text{L}$ arsenic). Children have higher levels of exposure, particularly if drinking water concentrations of arsenic are elevated, because of their smaller size and greater consumption of water relative to their size. Several state health departments and

public interest groups have expressed concern about children repeatedly exposed to arsenic from playing on arsenic-treated desk or play structures. Some exposure and associated risk calculations exceed EPA's acceptable risk levels. Arsenic exposure can also occur if arsenic treated wood is burned or from breathing sawdust from treated wood.

Biological Properties

Soluble inorganic arsenic compounds, such as arsenic trioxide, are readily absorbed from the intestine (80-90%). Organic arsenic compounds found in seafood are not well absorbed. Arsenic can also be absorbed through the lungs and skin. Most of the arsenic in the blood is bound to red blood cells. Once ingested, inorganic arsenic is biotransformed by the liver to a methylated form of arsenic and excreted in the urine with a half-life of 3 to 5 days. Arsenic is also excreted in the outer layer of skin cells and sweat. Arsenic binds to sulfhydryl-containing proteins and concentrates in the hair and fingernails. At higher levels of exposure, white bands, called Mees' lines, are visible in the nails.

Wipe off this glass three times.
There is arsenic in it.
I hear messages from God
through the fillings in my teeth.

Anne Sexton (1928–1974)

Health Effects

The acute effects of inorganic arsenic poisoning are well known from incidence of suicidal, homicidal and accidental poisonings. Ingestion of 70 to 180 mg of arsenic trioxide can be fatal, but initial effects may be delayed for several hours.

Symptoms following oral ingestion include constriction of the throat with difficulty in swallowing, severe intestinal pain, vomiting, diarrhea, muscle cramps, severe thirst, coma and death. If the patient survives the acute symptoms there is often peripheral nervous system damage.

They put arsenic in his meat
And stared aghast to watch him eat;
They poured strychnine in his cup
And shook to see him drink it up

A.E. (Alfred Edward) Housman (1859–1936)

The symptoms of chronic arsenic exposure are most often associated with contaminated drinking water. Early signs of arsenic exposure are garlic odor on the breath, excessive perspiration, muscle tenderness and weakness, and changes in skin pigmentation. More advanced symptoms include anemia, reduced sensation in the hand and feet from damage to the peripheral sensory system (stocking and glove syndrome), peripheral vascular disease, skin changes on palms and soles, and liver and kidney involvement. Changes in circulation can lead to gangrene of extremities, especially of the feet, which have been

referred to as blackfoot disease. Hyperpigmentation and hyperkeratosis of palms and soles occurs in 6 to 3 months with repeated ingestion of 0.4 mg/kg per day. Many of the symptoms are dose and time dependent. In other words, repeated low levels of exposure over an extend period of time can produce effects similar to a one time high level of exposure.

Arsenic cause both skin and lung cancer. Skin cancer was observed over 100 years ago in patients treated with arsenical compounds and lung cancer was seen in smelter workers who chronically inhaled arsenic dust. Although arsenic is an established human carcinogen, it has been difficult to confirm and study in animal models. Arsenic readily crosses the placenta, but there appears to be increased methylation of arsenic to its organic form, which reduces its toxicity to the fetus.

Reducing Exposure

The only supernatural agents, which can in any manner be allowed to us moderns, are ghosts; but of these I would advise an author to be extremely sparing. These are indeed, like arsenic, and other dangerous drugs in physic, to be used with the utmost caution

Henry Fielding, 1917

The toxicity of chronic exposure to arsenic is well established and the best recommendation is to avoid arsenic exposure entirely. The most common home exposure is from contaminated drinking water and arsenic-treated lumber. Certain areas of the country have higher levels of arsenic in water. The EPA has lowered arsenic drinking water standard to 10 ppb and required water providers until January 2006 to meet the new standard.

Avoid inhalation of sawdust from arsenic-treated lumber, and never burn any treated lumber or sawdust. Families with decks, play equipment, furniture, or other structures made with arsenic-treated lumber should take steps to reduce exposure, especially to children. Home uses of arsenic-treated lumber has been phased out in the United States, but it is estimated that approximately 60 billion board feet of arsenic-treated lumber are still in use in the United States as of 2002, about enough to cover half the state of California with a deck two inches thick. Several state agencies have recommended that treated lumber on which children may play should be coated periodically with paint or other sealer to reduce hand contact and subsequent ingestion of arsenic. Those who choose to remove arsenic-treated decks or other structures may want to test the soil underneath to see if levels exceed state standards. And always wash your hands after coming in contact with any arsenic treated product.

Regulatory Standards

EPA – Drinking water 10 $\mu\text{g/L}$ (10 ppb)

EPA – RfD - 0.3 $\mu\text{g/kg/day}$ (inorganic chronic exposure)

OSHA – Workplace air - 0.5 mg/m^3

ATSDR – MRL – 0.3 $\mu\text{g/kg/day}$ (chronic exposure)

Recommendation and Conclusions

Arsenic is an ancient and well-known hazard and, along with lead and mercury, is an important environmental contaminant. The inorganic form is far more toxic than organic arsenic, which is commonly found in seafood. Arsenic contaminated drinking water is a worldwide problem that affects millions of people. Human exposure also occurs from arsenic treated lumber.

The best recommendation is to avoid or reduce exposure to inorganic arsenic.

More Information and References

Slide Presentation

- A Small Dose of Arsenic presentation material and references online:
www.asmalldoseoftoxicology.org
Web site contains presentation material related to the health effects of arsenic.

European, Asian, and International Agencies

- World Health Organization (WHO). Online: < <https://www.who.int/news-room/fact-sheets/detail/arsenic> > (accessed: 16 September 2020).
WHO arsenic in drinking water fact sheet. Inorganic arsenic is very toxic, can contaminate drinking water.
- World Health Organization (WHO). Arsenic in Drinking Water and Resulting Arsenic Toxicity in India & Bangladesh. Online:
<https://www.who.int/ipcs/assessment/public_health/arsenic/en/> (accessed: 16 September 2020).
WHO report on arsenic in drinking water.

North American Agencies

- Health Canada – Arsenic in Drinking Water. Online: < <https://www.canada.ca/en/health-canada/services/healthy-living/your-health/environment/arsenic-drinking-water.html> > (accessed: 16 September 2020).
Health Canada provides information on the health effects arsenic in drinking water.
- U.S. Food and Drug Administration (FDA). “Arsenic in Food and Dietary Supplements” Online: < <https://www.fda.gov/food/metals-and-your-food/arsenic-food-and-dietary-supplements> > (accessed: 16 September 2020).
FDA Guidance Document for Arsenic in Food and Dietary Supplements.
- U.S. Environmental Protection Agency (EPA) - Arsenic Compounds. Online: < https://cfpub.epa.gov/si/si_public_record_report.cfm?Lab=ORD&count=10000&dirEntryId=42498&searchall=&showcriteria=2&simplesearch=0×type= html > (accessed: 16 September 2020).
EPA site has general information and research on arsenic.
- U.S. Environmental Protection Agency (EPA) – Integrated Risk Information System – Inorganic Arsenic. Online: <<https://www.epa.gov/iris/inorganic-arsenic-meetings-webinars#:~:text=The%20IRIS%20Program%20has%20been%20utilizing%20webinars%20in,the%20inorganic%20arsenic%20assessment%20can%20be%20found%20below%3A> > (accessed: 16 September 2020).
Site contains EPA’s risk assessment evaluation of inorganic arsenic.
- U.S. Environmental Protection Agency (EPA) – Toxics Release Inventory (TRI) Program. Online: <<http://www.epa.gov/tri/>> (accessed: 16 September 2020).
Site has information on arsenic release in the United States.
- U.S. ATSDR – Agency for Toxic Substance Disease Registry – Toxicology Profile Series Arsenic. Online: <<https://emergency.cdc.gov/agent/arsenic/>> (accessed: accessed: 16 September 2020).
- U.S. National Research Council (NRC) - Arsenic in Drinking Water: 2001 Update. Online: < http://www.nap.edu/catalog.php?record_id=10194 > (accessed: 16 September 2020).
The NRC report on arsenic can be accessed from the their web site.

- U.S. Geological Services (USGS). Online: < https://www.usgs.gov/mission-areas/water-resources/science/arsenic-and-drinking-water?qt-science_center_objects=0#qt-science_center_objects > (accessed: 16 September 2020).
Site contains a map of United States showing arsenic in water.

Non-Government Organizations

- SOS Arsenic Poisoning In Bangladesh / India. Online: <http://www.sos-arsenic.net/>. (accessed: 17 September 2020).
Information in English, German, Spanish, and French on arsenic poisoning in Bangladesh and India.

References

- Environmentally healthy homes and communities. Children's special vulnerabilities. (2001). *Am Nurse*, 33(6), 26-38; quiz 39-40.
- Hall, A. H. (2002). Chronic arsenic poisoning. *Toxicol Lett*, 128(1-3), 69-72.
- Jiang, J. Q. (2001). Removing arsenic from groundwater for the developing world--a review. *Water Sci Technol*, 44(6), 89-98.
- Liu, J., Zheng, B., Aposhian, H. V., Zhou, Y., Chen, M. L., Zhang, A., & Waalkes, M. P. (2002). Chronic arsenic poisoning from burning high-arsenic-containing coal in Guizhou, China. *Environ Health Perspect*, 110(2), 119-122.
- Pott, W. A., Benjamin, S. A., & Yang, R. S. (2001). Pharmacokinetics, metabolism, and carcinogenicity of arsenic. *Rev Environ Contam Toxicol*, 169, 165-214.
- Rahman, M. M., Chowdhury, U. K., Mukherjee, S. C., Mondal, B. K., Paul, K., Lodh, D., Biswas, B. K., Chanda, C. R., Basu, G. K., Saha, K. C., Roy, S., Das, R., Palit, S. K., Quamruzzaman, Q., & Chakraborti, D. (2001). Chronic arsenic toxicity in Bangladesh and West Bengal, India--a review and commentary. *J Toxicol Clin Toxicol*, 39(7), 683-700.
- Smith, A. H., Lingas, E. O., & Rahman, M. (2000). Contamination of drinking-water by arsenic in Bangladesh: a public health emergency. *Bull World Health Organ*, 78(9), 1093-1103.
- WHO. (2000). Towards an Assessment of Socioeconomic Impact of Arsenic Poisoning in Bangladesh. World Health Organization, Sustainable Development and Healthy Environments, WHO/SDE/WSH/00.4, 1-42.
- Yu, H. S., Lee, C. H., Jee, S. H., Ho, C. K., & Guo, Y. L. (2001). Environmental and occupational skin diseases in Taiwan. *J Dermatol*, 28(11), 628-631.

Information from Wikipedia (accessed: 17 September 2020)

- General overview - <https://en.wikipedia.org/wiki/Arsenic>

- Arsenic poisoning – map arsenic in ground water - https://en.wikipedia.org/wiki/Arsenic_poisoning
https://en.wikipedia.org/wiki/Arsenic_contamination_of_groundwater
- Arsenic biochemistry - https://en.wikipedia.org/wiki/Arsenic_biochemistry

New references

Carlin DJ, Naujokas MF, Bradham KD, Cowden J, Heacock M, Henry HF, Lee JS, Thomas DJ, Thompson C, Tokar EJ, Waalkes MP, Birnbaum LS, Suk WA. 2016. Arsenic and environmental health: state of the science and future research opportunities. *Environ Health Perspect* 124:890–899; <http://dx.doi.org/10.1289/ehp.1510209>

Helena Skroder Lovborn, Maria Kippler Ying Lu, Sultan Ahmed, Doris Kuehnelt, Rubhana Raqib, and Marie Vahter Arsenic Metabolism in Children Differs From That in Adults. *TOXICOLOGICAL SCIENCES*, 152(1), 2016, 29–39

The Challenge of Regulating Arsenic in Rice. *Environmental Health Perspectives* • volume 123 | number 1 | January 2015

Michael F. Hughes, Barbara D. Beck, Yu Chen, Ari S. Lewis, and David J. Thomas. Arsenic Exposure and Toxicology: A Historical Perspective. *TOXICOLOGICAL SCIENCES* 123(2), 305–332 (2011) doi:10.1093/toxsci/kfr184